

Application No.: 10/705,665


Docket No.: 21581-00240-US1

AMENDMENTS TO THE CLAIMS

In the claims:

Please amend the claims as follows:

1. (Twice Amended) A process for producing a 5-hydroxy-3-oxopentanoic acid derivative of the following formula (IV):



wherein  $R^1$  represents any of an alkyl group of 1 to 12 carbon atoms, an aryl group of 6 to 12 carbon atoms and an aralkyl group of 7 to 12 carbon atoms; and  $R^2$  represents any of hydrogen, an alkyl group of 1 to 12 carbon atoms which may have a substituent, an alkenyl group of 2 to 12 carbon atoms which may have a substituent, an aryl group of 6 to 12 carbon atoms which may have a substituent, a cyano group, a carboxyl group and an alkoxy carbonyl group,

which comprises [permitting] adding a lithium amide of the following formula (III):



wherein  $R^4$  and  $R^5$  may be the same or different and each represents any of an alkyl group of 1 to 12 carbon atoms, an aryl group of 6 to 12 carbon atoms, an aralkyl group of 7 to 12 carbon atoms and a silyl group,


to [act upon] a mixture of an acetic acid ester of the following formula (I) and a 3-hydroxypropionic acid derivative of the following formula (II) at a temperature not below  $-20^{\circ}\text{C}$  to conduct reaction:

Application No.: 10/705,665

Docket No.: 21581-00240-US1




wherein  $\text{R}^1$  represents any of an alkyl group of 1 to 12 carbon atoms, an aryl group of 6 to 12 carbon atoms and an aralkyl group of 7 to 12 carbon atoms:



wherein  $\text{R}^2$  represents any of hydrogen, an alkyl group of 1 to 12 carbon atoms which may have a substituent, an alkenyl group of 2 to 12 carbon atoms which may have a substituent, an aryl group of 6 to 12 carbon atoms which may have a substituent, an aralkyl group of 7 to 12 carbon atoms which may have a substituent, a cyano group, a carboxyl group and an alkoxycarbonyl group;  $\text{R}^3$  represents any of an alkyl group of 1 to 12 carbon atoms, an aryl group of 6 to 12 carbon atoms and an aralkyl group of 7 to 12 carbon atoms; and  $\text{R}^2$  and  $\text{R}^3$  may be joined to each other to form a ring, in the presence of a magnesium halide.

6. (Amended) A process for producing a 5-hydroxy-3-oxopentanoic acid derivative of the following formula (IV):



wherein  $\text{R}^1$  represents any of an alkyl group of 1 to 12 carbon atoms, an aryl group of 6 to 12 carbon atoms and an aralkyl group of 7 to 12 carbon atoms; and  $\text{R}^2$  represents any of hydrogen, an alkyl group of 1 to 12 carbon atoms which may have a substituent, an alkenyl group of 2 to 12 carbon atoms which may have a substituent, an aryl group of 6 to 12 carbon atoms which may have a substituent, an aralkyl group of 7 to


Application No.: 10/705,665

Docket No.: 21581-00240-US1

12 carbon atoms which may have a substituent, a cyano group, a carboxyl group and an alkoxycarbonyl group, which comprises treating a mixture of an acetic acid ester of the following formula (I) and a 3-hydroxypropionic acid derivative of the following formula (II):



wherein  $\text{R}^1$  represents any of an alkyl group of 1 to 12 carbon atoms, an aryl group of 6 to 12 carbon atoms and an aralkyl group of 7 to 12 carbon atoms:



wherein  $\text{R}^2$  represents any of hydrogen, an alkyl group of 1 to 12 carbon atoms which may have a substituent, an alkenyl group of 2 to 12 carbon atoms which may have a substituent, an aryl group of 6 to 12 carbon atoms which may have a substituent, an aralkyl group of 7 to 12 carbon atoms which may have a substituent, a cyano group, a carboxyl group and an alkoxycarbonyl group;  $\text{R}^3$  represents any of an alkyl group of 1 to 12 carbon atoms, aryl group of 6 to 12 carbon atoms and an aralkyl group of 7 to 12 carbon atoms; and  $\text{R}^2$  and  $\text{R}^3$  may be joined to each other to form a ring,

with a Grignard reagent of the following formula (V):




wherein  $\text{R}^6$  represents any of an alkyl group of 1 to 12 carbon atoms, an aryl group of 6 to 12 carbon atoms and an aralkyl group of 7 to 12 carbon atoms; and X represents halogen,

to prepare a mixture of a compound of the following formula (VI) and an acetic acid ester of the above formula (I):


Application No.: 10/705,665

Docket No.: 21581-00240-US1



(VI)


wherein  $\text{R}^2$  represents any of hydrogen, an alkyl group of 1 to 12 carbon atoms which may have a substituent, an alkenyl group of 2 to 12 carbon atoms which may have a substituent, an aryl group of 6 to 12 carbon atoms which may have a substituent, an aralkyl group of 7 to 12 carbon atoms which may have a substituent, a cyano group, a carboxyl group and an alkoxycarbonyl group;  $\text{R}^3$  represents any of an alkyl group of 1 to 12 carbon atoms, an aryl group of 6 to 12 carbon atoms and an aralkyl group of 7 to 12 carbon atoms;  $\text{R}^2$  and  $\text{R}^3$  may be joined to each other to form a ring; and X represents a halogen atom, and [permitting] adding a lithium amide of the following formula (III):



(III)

wherein  $\text{R}^4$  and  $\text{R}^5$  may be the same or different and each represents any of an alkyl group of 1 to 12 carbon atoms, an aryl group of 6 to 12 carbon atoms, an aralkyl group of 7 to 12 carbon atoms and a silyl group to [act upon] the mixture at a temperature not below  $-20^\circ \text{C}$  to conduct reaction.

10.(Amended) A process for producing a 5-hydroxy-3-oxopentanoic acid derivative of the following formula (IV):



(IV)

Application No.: 10/705,665

Docket No.: 21581-00240-US1

wherein  $R^1$  represents any of an alkyl group of 1 to 12 carbon atoms, an aryl group of 6 to 12 carbon atoms and an aralkyl group of 7 to 12 carbon atoms; and  $R^2$  represents any of hydrogen, an alkyl group of 1 to 12 carbon atoms which may have a substituent, an alkenyl group of 2 to 12 carbon atoms which may have a substituent, an aryl group of 6 to 12 carbon atoms which may have a substituent, an aralkyl group of 7 to 12 carbon atoms which may have a substituent, a cyano group, a carboxyl group and an alkoxycarbonyl group,

which comprises [permitting] adding a lithium amide of the following formula (III):



wherein  $R^4$  and  $R^5$  may be the same or different and each represents any of any alkyl group of 1 to 12 carbon atoms, an aryl group of 6 to 12 carbon atoms, an aralkyl group of 7 to 12 carbon atoms and a silyl group,

to [act upon] a mixture of an acetic acid ester of the following formula (I) and a compound of the following formula (VI) at a temperature not below  $-20^\circ C$  to conduct reaction:



wherein  $R^1$  represents any of an alkyl group of 1 to 2 carbon atoms, an aryl group of 6 to 12 carbon atoms and an aralkyl group of 7 to 12 carbon atoms:



Application No.: 10/705,665

Docket No.: 21581-00240-US1

wherein  $R^2$  represents any of hydrogen, an alkyl group of 1 to 12 carbon atoms which may have a substituent, an alkenyl group of 2 to 12 carbon atoms which may have a substituent, an aryl group of 6 to 12 carbon atoms which may have a substituent, an aralkyl group of 7 to 12 carbon atoms which may have a substituent, a cyano group, a carboxyl group and an alkoxy carbonyl group;  $R^3$  represents any of an alkyl group of 1 to 12 carbon atoms, an aryl group of 6 to 12 carbon atoms and an aralkyl group of 7 to 12 carbon atoms;  $R^2$  and  $R^3$  may be joined to each other form a ring; and X represents a halogen atom.

17.(amended) The process according to claim 1, wherein the [compound (II) or (VI)] the 3-hydroxypropionic acid derivative of the formula (II) or the 5-hydroxy-3-oxopentanoic acid derivative of the formula (IV) is optically active.

18. (amended) The process according to claim 2 wherein, referring to the acetic acid ester of the formula (I),  $R^1$  represents a tert-butyl group.